

University of Groningen

Multiple aspects of contact allergy

Dittmar, Daan

DOI:
[10.33612/diss.95667994](https://doi.org/10.33612/diss.95667994)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2019

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):
Dittmar, D. (2019). *Multiple aspects of contact allergy: immunology, patch test methodology and epidemiology*. [Thesis fully internal (DIV), University of Groningen]. University of Groningen.
<https://doi.org/10.33612/diss.95667994>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

stellingen behorende bij het proefschrift

Multiple Aspects of Contact Allergy

1. The likelihood of cross-elicitation reactions to 2-methyletoxy-*p*-phenylenediamine in *p*-phenylenediamine allergic individuals is associated with the strength of the original *p*-phenylenediamine patch test reaction. (*this thesis*)
2. The newly developed hair dye molecule 2-methyletoxy-*p*-phenylenediamine is not a safe alternative for *p*-phenylenediamine allergic consumers. (*this thesis*)
3. Patch testing with a pre-loaded patch tests such as the TRUE Test® gives more consistent results than patch testing with investigator-loaded patch tests. (*this thesis*)
4. For contact allergens with a steep dose-response curve, a small change in concentration can have a strong impact on diagnostic performance. (*this thesis*)
5. A definitive definition for polysensitisation in contact allergy should be decided on by the international research community in order to investigate susceptibility to contact allergy further. (*this thesis*)
6. As high rates of contact allergy to hydroperoxides of limonene and hydroperoxides of linalool have shown once again, we should always remain vigilant to new contact allergens – or existing ones not yet recognised as such. (*this thesis*)
7. The patch test concentration for hydroperoxides of limonene and hydroperoxides of linalool should be higher than the currently advised test concentration. (*this thesis*)
8. Patients should be advised to avoid contact allergens they reacted positively to, in order to evaluate whether the skin complaints disappear, but this advice should include a disclaimer that for weak positive reactions the threshold for an elicitation reaction might not be met during low grade exposure.
9. Clinical relevance depends too much on subjective information and patient participation to be considered a reliable outcome measure for research, and should be interpreted with the necessary caution.
10. Patch testing is less reliable as a diagnostic test in patients with atopic dermatitis.
11. Everything we hear is an opinion, not a fact. Everything we see is a perspective, not the truth. (Marcus Aurelius)